

IMMUNIZATION**DOSES REQUIRED TO START SCHOOL****(STUDENTS LESS THAN 5 YEARS OLD)**

Diphtheria, Tetanus, Pertussis (DTaP/DTP/DT/Tdap/Td)	<u>4 DOSES</u> -PLUS A BOOSTER AT 4YRS <u>IF</u> 4 th DOSE BEFORE AGE 4
POLIO	<u>3 DOSES</u> -PLUS A BOOSTER AT 4 YRS <u>IF</u> 3 RD DOSE BEFORE AGE 4
MMR	<u>1 DOSE</u> -PLUS A BOOSTER AT 4 YRS <u>OR</u> 1 MONTH AFTER 1 ST DOSE
HIB	<u>3 DOSES</u> -(3 RD dose after 12 months old) <u>OR</u> 1 HIB AT OR AFTER 15 MONTHS OLD
HEPATITIS B	<u>3 DOSES</u>
VARICELLA (CHICKEN POX)	<u>1 DOSE</u> - GIVEN ON OR AFTER 1 ST B-DAY
HEPATITIS A	<u>2 DOSES</u> - 1 ST DOSE MUST BE ON OR AFTER 1 ST B-DAY
PCV7	SEE PREVNAR SCHEDULE (*Note below)

KINDERGARTEN THRU 12TH GRADE**(5 YEARS AND OLDER)**

Diphtheria, Tetanus, Pertussis (DTaP/DTP/DT/Tdap/Td)	<u>5 DOSES</u> -UNLESS 4 TH DOSE GIVEN AFTER 4 TH B-DAY (FOR STUDENTS 7 YRS OR OLDER-3 DOSES REQUIRED IF LAST DOSE AFTER 4 TH BIRTHDAY) *Tdap BOOSTER REQUIRED FOR 7TH GRADE STUDENTS if 5 years since last DTP Vaccine <u>FOR 8TH-12TH GRADE: Tdap BOOSTER REQUIRED</u> When 10 years have passed since last DTP vaccine (Td acceptable substitute if Tdap medically contraindicated)
POLIO	<u>4 DOSES</u> -UNLESS 3 RD DOSE GIVEN AFTER 4 TH B-DAY
MEASLES, MUMPS, RUBELLA (MMR)	<u>2 DOSES</u> -1 ST DOSE MUST BE ON OR AFTER 1 ST B-DAY
HEPATITIS B	<u>3 DOSES</u>
VARICELLA (CHICKEN POX)	<u>2 DOSES</u>
HEPATITIS A	<u>2 DOSES</u> -(1 ST DOSE MUST BE ON OR AFTER 1 ST B-DAY) **REQUIRED FOR KINDERGARTEN THRU 9th GRADE ENTRY ONLY
MENINGOCOCCAL VACCINE	<u>1 DOSE</u> 7 TH Thru 12 TH GRADE STUDENTS ONLY

1

PNEUMOCOCCAL CONJUGATE VACCINE (PCV7) REQUIREMENTS

Age of Child (Months)	Vaccination History	Required Doses
2-6*	0 doses	3 doses, 2 months apart
	1 dose	2 doses, 2 months apart
	2 doses	1 dose, 2 months after the most recent dose
7-11*	0 doses	2 doses, 2 months apart
	1 or 2 doses	1 dose, 2 months after the most recent dose
12-23	0 doses	2 doses, ≥ 2 months apart
	1 or 2 doses administered before age 12 months	2 doses, ≥ 2 months apart
	1 dose administered on or after age 12 months	1 dose, ≥ 2 months after the most recent dose
	2 or 3 doses administered before age 12 months	1 dose, ≥ 2 months after the most recent dose
24-59 Healthy Children	Any incomplete schedule†	1 dose, > 2 months after most recent dose
Children at high risk¹	Any incomplete schedule†	2 doses separated by 2 months

*Children in these age groups will need a booster dose on or after 12 months of age.

†Any of the scenarios for children aged 2-6 (with a booster on or after 12 months of age), 7-11 (with a booster on or after 12 months of age), and 12-23 months constitute a complete series.

¹Children with certain chronic conditions or immuno-supression conditions will also require a dose of pneumococcal polysaccharide vaccine (PPV23) in addition to PCV7 two months after the last PCV7.